Overview

Aims and Scope

Perspectives in Psychiatric Care (PPC) is recognized and respected as THE journal for advanced practice psychiatric nurses. The journal provides advanced practice nurses with current research, clinical application, and knowledge about psychiatric nursing, prescriptive treatment, and education. It publishes peer-reviewed papers that reflect clinical practice issues, psychobiological information, and integrative perspectives that are evidence-based. *Perspectives in Psychiatric Care* includes regular columns on the biology of mental illness and pharmacology, the art of prescribing, integrative perspectives, and private practice issues.

Readership

National and international, serving psychiatric/advanced practice mental health nurses and other mental health professionals

Keywords

psychiatry, mental health, nursing, psychiatric nursing, psychopharmacology, perspectives, care

Abstracting and Indexing Information

- Academic Search (EBSCO Publishing)
- Academic Search Alumni Edition (EBSCO Publishing)
- Academic Search Elite (EBSCO Publishing)
- Academic Search Premier (EBSCO Publishing)
- AgeLine Database (EBSCO Publishing)
- British Nursing Database (ProQuest)
- CINAHL: Cumulative Index to Nursing & Allied Health Literature (EBSCO Publishing)
- Current Contents: Social & Behavioral Sciences (Clarivate Analytics)
- Expanded Academic ASAP (GALE Cengage)
- Health & Medical Collection (ProQuest)
- Health Research Premium Collection (ProQuest)
- Health Source Nursing/Academic (EBSCO Publishing)
- Hospital Premium Collection (ProQuest)
- InfoTrac (GALE Cengage)
- Journal Citation Reports/Science Edition (Clarivate Analytics)
- Journal Citation Reports/Social Science Edition (Clarivate Analytics)
- Medical Database (ProQuest)
- MEDLINE/PubMed (NLM)
- Nursing & Allied Health Database (ProQuest)

Perspectives in PSYCHIATRIC CARE








Subscribe to this journal

More from this journal

- News
- Professional Opportunities



DIVERSITY

in Research Jobs

Please <u>contact us</u> to see your job listed here

Seasonal RN Registered Nurse Labor and Delivery

Fort Worth, TX





HOME ABOUT ✓

CONTRIBUTE V

BROWSE V

nescaren cibrary Frep (Froquest)

• Science Citation Index Expanded (Clarivate Analytics)

- SCOPUS (Elsevier)
- Social Sciences Citation Index (Clarivate Analytics)
- STEM Database (ProQuest)

Clinical Faculty, the Instructor, supports the University in fulfillment of its mission, purpose and go...

Wiley Online Library

Search



Login / Register

Editorial Board

Editor

Evelyn Parrish, PhD, PMHNP-BC

Associate Professor, University of Kentucky College of Nursing

Founding Editor Alice Clarke, RN, MA

Hillsdale, New Jersey, USA

Managing Editor

Rachel Yehl Email: ppc-admin@wiley.com

Editorial Board

Deborah Antai-Otong, MS, PMHCNS-BC

Mental Health Provider, VA North Texas Health Care System (VANTHCS), Dallas, TX, USA

Paul T. Clements, PhD, APRN, BC, CGS, DF-IAFN

Associate Clinical Professor, Drexel University, Philadelphia, PA, USA

Margaret England, PhD, RN

Associate Professor, University of Windsor, Windsor, ON, Canada

Cheryl Forchuk, PhD, RN

Professor, Nursing & Psychiatry, University of Western Ontario, Canada Assistant Director and Scientist, Lawson Health Research Institute, Canada

Peter Goossens, PhD, PMH-CNS, RN, FEANS

Professor in Mental Health, Saxion University of Applied Sciences Deventer, the Netherlands Nurse practitioner/Senior researcher, Dimence Mental Health, Deventer, the Netherlands; Senior Researcher, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands; Fellow, European Academy of Nursing Science

Cynthia Handrup, DNP, APRN, PMHCNS-BC

Clinical Assistant Professor, University of Illinois - Chicago, Chicago, IL, USA

Brenda Happell, RN, RPN, BA(Hons), Dip Ed., BEd., MEd, PhD

Professor of Contemporary Nursing, Faculty of Sciences, Engineering and Health, CQUniversity Australia

Vicki Hines-Martin, PhD, CNS, RN, FAAN

Associate Professor, University of Louisville School of Nursing, Louisville, KY, USA

Norman Keltner, EdD, APRN-BC

Professor, University of Alabama at Birmingham, Birmingham, Alabama, USA

Berno van Meijel, RN, PhD

Submit an Article


Browse free sample issue

Get content alerts

Subscribe to this journal

More from this journal

- News
- Professional Opportunities





in Research Jobs

Please contact us to see your job listed here

Seasonal RN Registered Nurse Labor and Delivery

Fort Worth, TX

Job Description Description SHIFT: PRN SCHEDULE: PRN/Per Diem Seasonal Program - RN Registered NurseLabor and Delivery / L&D Seasonal PRN Program R...

Employer: Medical City Alliance

Apply for this job

Perspectives in PSYCHIATRIC CARE

HOME

ABOUT ~

CONTRIBUTE ~

BROWSE V





Diane Snow, PhD, RN, PMHNP-BC, CARN


Clinical Professor and Director, Psychiatric-Mental Health Nurse Practitioner Program, University of Texas at Arlington School of Nursing, Arlington, TX, USA

Roberta Waite, EdD, PMHCNS-BC, FAAN

Associate Professor and Assistant Dean of Academic Integration of Community Programs and Doctoral Nursing Department, Drexel University, Philadelphia, PA, USA

Mary Weber, PhD, APRN, PMHNP-BC

Endowed Professor in Psychiatric Nursing, University of Colorado College of Nursing, Aurora, CO, USA





Master Journal List Search Journals

Match Manuscript

Downloads Help Center

Welcome, Zehra Çalışkan

⇔ Settings **→** Log Out

NEW

Clear All

V

The power of the Web of Science™ on your mobile device, wherever inspiration strikes.

Dismiss

Learn More

Already have a manuscript?

Use our Manuscript Matcher to find the best relevant journals!

Find a Match

efine Your Search Results

Perspectives in Psychiatric Care

rt By: Relevancy •

earch Results

xact Match Found

ound 893 results (Page 1)

Share These Results

Web of Science Coverage

Filters

Open Access 🔒 🔍



Country / Region

Language

PERSPECTIVES IN PSYCHIATRIC CARE

Publisher: WILEY, 111 RIVER ST, HOBOKEN, USA, NJ, 07030-5774

ISSN / eISSN: 0031-5990 / 1744-6163

Web of Science Core Collection: Science Citation Index Expanded | Social Sciences Citation Index

Additional Web of Science Indexes: Current Contents Social And Behavioral Sciences | Essential Science Indicators

Search



PERSPECTIVES IN PSYCHIATRIC CARE

ISSN

0031-5990

EISSN

1744-6163

JCR ABBREVIATION

PERSPECT PSYCHIATR C

ISO ABBREVIATION

Perspect. Psychiatr. Care

Journal information

EDITION

Social Sciences Citation Index (SSCI)

CATEGORY

PSYCHIATRY - SSCI

NURSING - SSCI

Science Citation Index Expanded (SCIE)

NURSING - SCIE

PSYCHIATRY - SCIE

LANGUAGES REGION 1ST ELECTRONIC JCR YEAR

English USA 1997

Publisher information

PUBLISHER

ADDRESS PUBLICATION FREQUENCY

WILEY 111 RIVER ST, HOBOKEN 07030-

5774, NJ

<u>◆ Export</u> Customize

4 issues/year

Year	Total Citations	Journal impact factor	JIF without self cites	5 Year Impact Factor	Immediacy Index	Citable items	% of articles in Citable items	Average JIF Percentile
2020	1,276	2.186	1.904	2.185	0.424	290	96.90	45.922
2019	721	1.273	1.116	1.417	0.316	98	92.86	33.245
2018	661	1.240	1.147	1.374	0.405	79	91.14	34.212
2017	494	1.015	1.000	1.036	0.452	42	92.86	28.708
2016	501	1.130	1.087	1.205	0.212	33	93.94	34.453
2015	425	1.000	0.857	1.213	0.382	34	91.18	34.529
2014	338	0.651	0.619	1.000	0.086	35	94.29	18.797
2013	334	0.712	0.712	1.242	0.171	35	91.43	22.694
2012	293	1.038	0.981	1.242	0.143	28	92.86	46.329
2011	296	1.298	1.246	1.387	0.292	24	100.00	61.388
2010	234	1.058	0.981	1.144	0.286	28	96.43	46.454
2009	207	1.000	0.884	1.195	0.172	29	100.00	44.817
2008	211	1.103	0.846	1.207	0.478	23	91.30	49.362
2007	153	1.062	0.781	0.937	0.250	20	100.00	47.096
2006	139	0.800	0.600	n/a	0.316	19	100.00	26.878
2005	90	0.067	0.033	n/a	0.231	13	92.31	1.974

2003 75 0.333 0.200 n/a 0.077 13 100.00 11.79 2002 96 0.240 0.200 n/a 0.000 17 94.12 10.50 2001 80 0.526 0.263 n/a 0.000 13 92.31 40.60 2000 103 0.545 0.455 n/a 0.083 12 100.00 40.30									
2002 96 0.240 0.200 n/a 0.000 17 94.12 10.5 2001 80 0.526 0.263 n/a 0.000 13 92.31 40.6 2000 103 0.545 0.455 n/a 0.083 12 100.00 40.3	2004	91	0.571	0.457	n/a	0.000	12	100.00	28.811
2001 80 0.526 0.263 n/a 0.000 13 92.31 40.63 2000 103 0.545 0.455 n/a 0.083 12 100.00 40.34	2003	75	0.333	0.200	n/a	0.077	13	100.00	11.793
2000 103 0.545 0.455 n/a 0.083 12 100.00 40.34	2002	96	0.240	0.200	n/a	0.000	17	94.12	10.571
· · · · · · · · · · · · · · · · · · ·	2001	80	0.526	0.263	n/a	0.000	13	92.31	40.639
1999 99 0.267 0.233 n/a 0.000 7 100.00 19.0	2000	103	0.545	0.455	n/a	0.083	12	100.00	40.347
	1999	99	0.267	0.233	n/a	0.000	7	100.00	19.070
1998 90 0.400 0.229 n/a 0.000 15 100.00 38.73	1998	90	0.400	0.229	n/a	0.000	15	100.00	38.720
1997 15 0.111 0.111 n/a 0.000 15 100.00 10.1-	1997	15	0.111	0.111	n/a	0.000	15	100.00	10.145

Accelerating innovation © 2021 Clarivate Copyright notice Terms of Use Privacy statement Cookie policy

Perspectives in

PSYCHIATRIC CARE

The Journal for Advanced Practice Psychiatric Nurses



Perspectives in PSYCHIATRIC CARE

The Journal for Advanced Practice Psychiatric Nurses

Volume 57, Number 3, July 2021

				•
$-\epsilon$	ır	tı	٦r	ia

979 The ongoing conundrum of long-term benzodiazepine use: What our patients need to know Evelyn Parrish

Original Articles

- 980 Effect of acupressure on agitation in the elderly with dementia who receive institutional care: A pilot study
 Serap Bayram, Özlem Altınbaş Akkaş, and Esra Usta
- 789 The hopes of nurses who care for people diagnosed with chronic mental illness in closed wards Kyung Mi Sung, Sun Ah Park, Jin Hee Ham
- 1000 Relationship between the social support and psychological resilience levels perceived by nurses during the COVID-19 pandemic: A study from Turkey

 Tülay Kılınc and Aslı Sis Celik
- 1009 Self-compassion and nursing competency among Japanese psychiatric nurses Yusuke Kurebayashi
- 1019 Quality of sexual life and factors affecting it in married women undergoing depression treatment
 Alev Ekemen and Kerime Derya Beydağ
- 1026 Relationship between nomophobia, fear of missing out, and perceived work overload in nurses in Turkey Haydar Hoşgör, Filiz Coşkun, Figen Çalişkan, and Derya Gündüz Hoşgör
- 1034 Determination of the relationship between the beliefs of nursing students toward mental illnesses and their empathic tendency levels

 Hülya Kök Eren and Nermin Gürhan
- 1041 The effect of coronavirus (COVID-19) outbreak on the mental well-being and mental health of individuals Adem Sümen and Derya Adibelli
- 1052 Preservice teachers' nomophobia levels, sense of loneliness and adjustment to college life Yadigar Çevik-Durmaz, Eylem Yalçinkaya-Önder, and Serkan Timur
- 1073 Risk factors and adverse clinical outcomes of postoperative delirium in Thai elderly patients:

 A prospective cohort study

Tanyong Pipanmekaporn, Yodying Punjasawadwong, Nahathai Wongpakaran, Tinakon Wongpakaran, Kittinan Suwannachai, Kaweesak Chittawatanarat, and Sirirat Mueankwan

- 1083 Self-esteem and sexual quality of life among obese women Hilal Türkben Polat and Emine Kaplan Serin
- 1088 State anxiety levels of nurses providing care to patients with COVID-19 in Turkey Emel Bahadir-Yilmaz and Arzu Yüksel

1095	Effect of progressive muscle relaxation exercise on clinical stress and burnout in student nurse interns Sıdıka Pelit-Aksu, Sultan Özkan-Şat, Şengül Yaman-Sözbir, and Ayten Şentürk-Erenel
1103	An investigation of individuals' health anxiety during the COVID-19 pandemic within the framework of the functional health patterns Sultan Alan, Sule Gokyildiz Surucu, Burcu Avcibay Vurgec, and Ayseren Cevik
1114	Examination of nursing students' anxiety levels related to clinical practice with respect to peer support Hatice Tambağ
1120	The effects of communication skills on resilience in undergraduate nursing students in Turkey Serap Yıldırım, Ebru Kazandı, Kadriye Cirit, and Hatice Yağız
1126	"I'm a hero, but": An evaluation of depression, anxiety, and stress levels of frontline healthcare professionals during COVID-19 pandemic in Turkey Handan Alan, Feride Eskin Bacaksiz, Hanife Tiryaki Sen, Gulcan Taskiran Eskici, Emel Gumus, and Arzu K. Harmanci Seren
1137	The effect of maternal depression symptoms on the outcomes of infant care Salih Güler and Arzu Akcan
1145	Practice change: Social Media Screening Questionnaire to identify high-risk adult psychiatric patients Cara A. Lutzow, Grace Hubbard, Cheryl Giscombe, and Lawrence Greenberg
1150	Depression and suicidality among Bangladeshi students: Subject selection reasons and learning environment as potential risk factors Najmuj Sakib, Merajul Islam, Md. Syeed Al Habib, A. K. M. Israfil Bhuiyan, Md. Mahbub Alam, Noshin Tasneem, Moazzem Hossain, Sheikh Mohammed Shariful Islam, Mark D. Griffiths, and Mohammed A. Mamun
1163	Psychiatric inpatients' view of the therapeutic environment: A Turkish interview study Işıl Işık, Gül Ergün, Gül Dikeç, Volkan Ayaz, and Bülent Kadri Gültekin
1175	The effect of interpersonal relational role analysis on nursing students' anxiety levels and interpersonal problem-solving orientation

Veysel Kaplan and Gülsüm Ançel

- Effects of a nonviolent communication-based training program for inpatient alcoholics in South Korea Jeongwoon Yang and Sungjae Kim
- 1195 Secondary traumatic stress among midwifery students Fadime Bayri Bingol, Meltem Demirgoz Bal, Merve Aygun, and Edanur Bilgic
- 1202 Relationships between perceived support types and the job satisfaction levels of nurses Şehrinaz Polat, and Banu Terzi
- 1212 Knowledge, attitudes, and practices toward the coronavirus disease 2019 (COVID-19) among nurses in Lebanon

Danielle Saadeh, Hala Sacre, Souheil Hallit, Rita Farah, and Pascale Salameh

1222 Experiences and opinions of nurses working in psychiatric wards on assessing patients' sexual health: A qualitative study

Emre Ciydem, Ozlem Kackin, and F. Yasemin Kutlu

1231 Burnout levels and sleep quality of COVID-19 heroes

Aylin Aydin Sayilan, Nurşen Kulakaç, Sevda Uzun

1237 Monitoring for the metabolic side effects of second-generation antipsychotic medications: Psychiatrists' views and practices

Bayan Abdulhaq, Latefa A. Dardas, and Omar Sami

1244 Bipolar disorder: Treatment strategies for women of childbearing age Brayden N. Kameg

1250	The effect of progressive muscle relaxation exercises on compassion satisfaction, burnout, and compassion fatigue of nurse managers Remziye Semerci, Gülnur Öztürk, Melahat Akgün Kostak, Saadet Elmas, Ali İhsan Danacı, and Sadber Musbeg
1257	The effect of shyness levels of women on early diagnosis attitudes towards cervical cancer Safiye Ağapinar Şahin, and Mine Bekar
1266	Body perception and sexuality of bariatric surgery patients Sibel Çaynak, Nur E. Boyacıoğlu, and Münire Temel
1273	The effectiveness of Tai Chi Chuan on fear of movement, prevention of falls, physical activity, and cognitive status in older adults with mild cognitive impairment: A randomized controlled trial Canan Birimoglu Okuyan and Ebru Deveci
1282	The effect of revised process recording on self-focus and self-compassion: A randomized controlled pilot study Yusuke Kurebayashi
1292	Turkish validity and reliability of the Internalized AIDS-Related Stigma Scale Servet Öztürk, Çağlar Şimşek, Elif Ardıç, and Canan Ağalar
1299	Attitudes of students studying in various fields related to health services toward gender roles and intimate partner violence Funda Evcili and Gulseren Daglar
1305	Experiences of Dutch ambulance nurses in emergency care for patients with acute manic and/or psychotic symptoms: A qualitative study Thea H. Daggenvoorde, Josette M. van Klaren, Harm J. Gijsman, Hester Vermeulen, and Peter J. J. Goossens
1313	Factors associated with problematic social media use among a sample of Lebanese adults: The mediating role of emotional intelligence Sam Barbar, Chadia Haddad, Hala Sacre, Dina Dagher, Marwan Akel, Nelly Kheir, Pascale Salameh, Souheil Hallit, and Sahar Obeid
1323	Caregivers' psychological suffering and posttraumatic growth after patient death Eunmi Lee and Yujeong Kim
1331	Association between depression or sleep problems and working time mismatch among Korean workers Eunmi Lee, Yujeong Kim, and Haeyoung Lee
1340	Nurses' views and applications on palliative care Diğdem Lafci, Ebru Yildiz, and Seda Pehlivan
1347	Factors associated with overweight and obesity in Lebanese male patients with schizophrenia Maha Kassem, Chadia Haddad, Christelle Daccache, Claire Hayek, Souheil Hallit, and Francois Kazour
1356	Levels of fear and uncertainty regarding the spread of coronavirus disease (COVID-19) among university students Nadia B. Elsharkawy and Enas M. Abdelaziz
1365	Locked versus open ward environments and restrictive measures in acute psychiatry in Greece: Nursing students' attitudes and experiences Evdokia Missouridou, Afroditi Zartaloudi, Chrisoula Dafogianni, John Koutelekos, Evangelos Dousis, Eugenia Vlachou, and Eleni Evagelou
1376	The psychological impact of the Syrian crisis on refugees living in Lebanon Mohamed Hendaus, Nisreen Mourad, Samar Younes, Dalal Hammoudi, Mohamad Rahal, and Iman Basheti
1382	Developing and psychometric properties of the caring behaviors assessment tool-patient version-short form Mehtap Akgün, Selma Turan Kavradim, İlkay Boz, Zeynep Özer, and Sherill Nones Cronin
1390	Impact of a video or photography-assisted adaptation program on the concerns and social stigmatization-related perceptions of first-time psychiatric inpatients Halil İbrahim Bilkay and Oya Sevcan Orak

1399	Psychological factors associated with the spread of Coronavirus disease 2019 (COVID-19) among nurses working in health sectors in Saudi Arabia Hana M. Abu-Snieneh
1409	Evaluation of the chronic disease management and depression levels of people over 65 years of age during the COVID-19 pandemic period Aylin Y. Irmak, Ülfiye Çelikkalp, and Galip Ekuklu
1417	Nurses' perceptions toward factors that cause medication errors in Jordan: A qualitative study Luma Ali, Ahmad Saifan, Intima Alrimawi, and Maha Atout
1425	Examination of evidence-based nursing attitudes and the relationship with professional self-concept in nurses working in a psychiatric hospital Sibel Asi Karakaş, Özlem Şahin Altun, Hatice Polat, and Zeynep Öztürk
1433	Psychological distress and mental health literacy in university students Şenay Pehlivan, Makbule Tokur Kesgin, and Pelin Uymaz
1442	The relationship between nursing students' technology addiction levels and attitudes toward e-learning during the COVID-19 pandemic: A cross-sectional study Nur Guven Ozdemir and Munevver Sonmez
1449	Anxiety, depression, and knowledge level in postpartum women during the COVID-19 pandemic Gulten Guvenc, İlknur Yesilcinar, Fulden Ozkececi, Emine Öksüz, Coşkun F. Ozkececi, Dilek Konukbay, Gulsah Kok, and Kazım E. Karasahin
1459	The Compassion Fatigue-Short Scale for healthcare professionals: A Turkish study of validity and reliability Serap Yıldırım and Funda Cavcav
1466	The palliative care difficulty levels of the nursing students during clinical practices Seher Ünver, Zeynep Kızılcık Özkan, and Beyza N. Girgin
1474	Resilience and mindfulness in nurse training on an undergraduate curriculum Andrew E. P. Mitchell
1482	Identification of the relationship between nicotine dependence levels and cessation fatigue Ayse Berivan Bakan, Betül Aktas, and Gülpınar Aslan
1489	Professionals' views on patient involvement in acute psychiatric wards: A qualitative study Tiina Kortteisto, Minna Laitila, and Anneli Pitkänen
1497	Sexual myths and attitudes regarding sexuality of nursing students: A mixed method study Ayse Akalin, Birgul Ozkan
1505	Determination of humant life catisfaction, and stress coning styles of nevents with dischlad shildren
1203	Determination of burnout, life satisfaction, and stress coping styles of parents with disabled children Zehra Çalışkan, Derya Evgin, Nuray Caner, Bahriye Kaplan, and Gonca Özyurt
1515	Intimate partner violence during pregnancy in Turkey: A systematic review and meta-analysis

Case Reports

1528 The first COVID-19 triadic (homicide!)-suicide pact: Do economic distress, disability, sickness, and treatment negligence matter?

Mohammed A. Mamun

Nur Elçin Boyacioğlu, Sevil Günaydın, Neslihan Keser Özcan, and Hüsniye Dinç Kaya

1532 Factitious disorder, presenting haemolacria in a Pakistani tribal illiterate female: A rare case report Abdul Wali Khan, Asad Ali Khan, Muhammad Ishaq, Irfan Ullah, and Mohammed A. Mamun

Letter to the Editor

1536 Impact of daytime routine modifications on people with severe intellectual disability amid COVID-19 pandemic

Luigi Vetri, Maurizio Elia, Girolamo Aurelio Vitello, Donatella Greco, Catalda Gagliano, Maria Cristina Costanzo, Giusi Romeo, and Sebastiano Antonino Musumeci

Information for Authors

Perspectives in Psychiatric Care (PPC) is the peer-reviewed, international interdisciplinary resource for advanced practice psychiatric nurses and for nurse psychotherapists. The journal publishes original articles about innovations in clinical practice. education, or administration; case presentations; public policy information; synthesis of research findings; and original research. Contributions should (a) clearly specify the nursing problems/issues, (b) make a compelling argument for the way in which the author addresses the issues, (c) provide information that will facilitate clinical judgments, (d) describe how clinicians can best use the information presented, and (e), where applicable, include a list of additional resources (including Web sites) that provide related information for clinical judgment.

Manuscript submission

PPC accepts original contributions through online submission to ScholarOne Manuscripts at http://mc.manuscriptcentral.com/ppc.

Please log in to the site with your username and password and choose the "Author Center" option. (If you are a new contributor, first choose the "Create an Account" tab in the top right corner of the window.)
Complete instructions are provided to guide you through the submission process, but questions about this process can be directed to the Editorial Office at ppc-admin@wiley.com

Prepare separate files for the (a) title page, (b) abstract, (c) body and references, (d) figures, (e) tables, and (f) pictures. Contributions should be typed in 12-point font, double-spaced, with 1-inch margins, and no longer than 20 pages total. Identify each page with a number in the upper right corner.

The submitting author should enter his/ her contact details, as well as information about the co-authors. To ensure a double-blind peer-review process, authors' names and other identifying information should not appear in any uploaded files. The title page should be uploaded separately from the other files and designated as a "supplementary file NOT for review."

Include a structured abstract of no more than 100 words. Headings for research abstracts are Purpose, Design and Methods, Findings, and Practice Implications. For other abstract types, please use Purpose, Conclusions, and Practice Implications. Include 3 to 5 relevant keywords as search terms.

Contributions should conclude with a final paragraph under the heading *Implications for Nursing Practice*. Authors should address this issue to help readers judge the existing evidence for decision-making.

Each figure and table should be referenced within the text and include a short legend.

References

Whenever possible, references should be limited to publications within the last 5 years. References should follow the format of the Publication Manual of the American Psychological Association, 6th ed. (2010). For example:

Journal article: Pearson, G. S. (2009). When to

Pearson, G. S. (2009). When to forgive [Editorial]. Perspectives in Psychiatric Care, 45(3), 159-160. doi: 10.1111/j.1744-6163.2009.00217.x

Advance online publication: May, K. M., & Rew, L. (2009). Mexican American youths' and mothers' explanatory models of diabetes prevention. *Journal for Specialists in Pediatric Nursing*. Advance online publication. doi: 10.1111/j.1744-6155.2009.00207.x

Chapter in a book:

Bryant, R. (2004). In the aftermath of trauma: Normative reactions and early interventions. In G. Rosen (Ed.), Posttraumatic stress disorder: Issues and controversies (pp. 187–211). West Sussex, UK: John Wiley & Sons.

Reference with more than 7 authors: Bramley, D., Riddell, T., Whittaker, R., Corbett, T., Lin, R-B., Wills, M., ... Rodgers, A. (2005). Smoking cessation using mobile phone text messaging is as effective in Maori as non-Maori. New Zealand Medical Journal, 118(1216), 1491.

General Formatting Requirements Follow the style of the Publication Manual of the American Psychological Association, 6th ed. (2010). Authors can review the changes between the 5th ed. and 6th ed. of the manual at: http://www.apastyle.org. Use generic names for medications. Put all measurements in the metric system. Acknowledge any funding (monetary or in-kind support or provision of equipment) for conduct of the research or preparation of the manuscript. If any author has a financial stake in the funding agency or a product mentioned in the manuscript, state this also.

Art work, figures, charts, tables
Authors are responsible for
obtaining permission to reproduce
any copyrighted material, including
text, tables, photographs, and
figures; see Copyright section below.
If photographs of people are used
and the person is identifiable, copies
of the person's permission to use the
photograph must accompany the

Label and title all figures and include the necessary legend(s). Submit vector graphics or line art saved as Encapsulated PostScript (EPS) or PDFs and halftones or photographic images saved as Tagged Image Format Files (TIFF). Please find detailed information online at http:// authorservices.wiley.com/bauthor/ illustration.asp for electronic submissions. Do NOT submit figures as Word (.doc), Excel (.xls), or PowerPoint (.ppt) files.

Refereed publication

All manuscripts will be reviewed by the Editor, and if deemed appropriate for PPC, at least two members of the review panel. PPC uses a double-blind peer-review process. Accepted manuscripts may be revised to conform to the journal's standards and style.

Copyright

If your paper is accepted, the author identified as the formal corresponding author for the paper will receive an email prompting them to login into Author Services; where via the Wiley Author Licensing Service (WALS) they will be able to complete the license agreement on behalf of all authors on the paper.

For authors signing the copyright transfer agreement

If the OnlineOpen option is not selected the corresponding author will be presented with the copyright transfer agreement (CTA) to sign. The terms and conditions of the CTA can be previewed in the samples associated with the Copyright FAQs below:

CTA Terms and Conditions http://authorservices.wiley.com/ bauthor/faqs_copyright.asp

For authors choosing OnlineOpen If the OnlineOpen option is selected the corresponding author will have a choice of the following Creative Commons License Open Access Agreements (OAA):

Creative Commons Attribution License OAA Creative Commons Attribution Non-Commercial License OAA Creative Commons Attribution Non-Commercial-NoDerivs License OAA

To preview the terms and conditions of these open access agreements please visit the Copyright FAQs hosted on Wiley Author Services http://authorservices.wiley.com/bauthor/faqs_copyright.asp and visit http://www.wileyopenaccess.com/details/content/12f25db4c87/Copyright-License.html.

If you select the OnlineOpen option and your research is funded by The Wellcome Trust and members of the Research Councils UK (RCUK) you will be given the opportunity to publish your article under a CC-BY license supporting you in complying with Wellcome Trust and Research Councils UK requirements. For more information on this policy and the Journal's compliant self-archiving policy please visit: http://www.wiley.com/go/funderstatement.

NIH Public Access Mandate Authors can note on the CTA that they received NIH funding, and Wiley Blackwell will post the accepted version of the manuscript to PubMed Central on the author's behalf. For more information, please go to: http://www.wiley.com/go/nihmandate

Author order

After a manuscript has been submitted, any change to the listed order of the authors must be accompanied by a statement indicating the change and signatures from each of the authors signaling their agreement. This statement must be sent to the Managing Editor by email (ppc-admin@wiley.com)

Dual Submission

It is unethical to submit a manuscript to more than one journal at a time. If the manuscript was submitted to Journal A and received a decision of "major revision," for example, after the first round of review, the manuscript can not be submitted to Journal B until the author has formally requested that the manuscript be "withdrawn" from consideration from Journal A. (Receiving a decision of "reject" ends the submission process and the manuscript may be submitted elsewhere.)

Letters to the Editor

Letters to the Editor are welcomed and will be published on a space-available basis.

Letters may be revised to conform to the journal's standards and style. Please send correspondence to Evelyn Parrish, PhD, APRN-BC, at Evelyn.Parrish@uky.edu

Authors for whom English is a second language may choose to have their manuscript professionally edited before submission to improve the English. Information about Wiley's editing service can be found at http://wileyeditingservices. com/en/. All services are paid for and arranged by the author, and use of one of these services does not guarantee acceptance or preference for publication.

Online production tracking via Author Services

Author Services enables an author to track his/her article—after it has been accepted—through the production process to publication online and in print. An author can check an article's status online and choose to receive automated e-mails at key stages of production. For more information, visit http://authorservices.wiley.com/bauthor/.

Offprints

Authors will have access to a free PDF of their published article via Wiley Blackwell's Author Services. Information will be provided during the production process on registering for this service.

Perspectives in PSYCHIATRIC CARE

Editorial Board

Editor

Evelyn Parrish, PhD, PMHNP-BC

Associate Professor, University of Kentucky College of Nursing

Founding Editor

Alice Clarke, RN, MA

Hillsdale, New Jersey, USA

Managing Editor

Rachel Yehl

Email: ppc-admin@wiley.com

Editorial Board

Deborah Antai-Otong, MS, PMHCNS-BC

Mental Health Provider, VA North Texas Health Care System (VANTHCS), Dallas, TX, USA

Paul T. Clements, PhD, APRN, BC, CGS, DF-IAFN

Associate Clinical Professor, Drexel University, Philadelphia, PA, USA

Margaret England, PhD, RN

Associate Professor, University of Windsor, Windsor, ON, Canada

Cheryl Forchuk, PhD, RN

Professor, Nursing & Psychiatry, University of Western Ontario, Canada

Assistant Director and Scientist, Lawson Health Research Institute, Canada

Peter Goossens, PhD, PMH-CNS, RN, FEANS

Professor in Mental Health, Saxion University of Applied Sciences Deventer, the Netherlands Nurse practitioner/Senior researcher, Dimence Mental Health, Deventer, the Netherlands; Senior Researcher, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands; Fellow, European Academy of Nursing Science

Cynthia Handrup, DNP, APRN, PMHCNS-BC

Clinical Assistant Professor, University of Illinois - Chicago, Chicago, IL, USA

Brenda Happell, RN, RPN, BA(Hons), Dip Ed., BEd., MEd, PhD

Professor of Contemporary Nursing, Faculty of Sciences, Engineering and Health, CQUniversity Australia

Vicki Hines-Martin, PhD, CNS, RN, FAAN

Associate Professor, University of Louisville School of Nursing, Louisville, KY, USA

Norman Keltner, EdD, APRN-BC

Professor, University of Alabama at Birmingham, Birmingham, Alabama, USA

Berno van Meijel, RN, PhD

Professor of Mental Health Nursing, Inholland University of Applied Sciences (Amsterdam), VU University Medical Center and Parnassia Bavo Psychiatric Institute (The Hague), The Netherlands

Karen M. Ragaisis MSN, APRN, PMHNP-BC, PMHCNS-BC, CARN

Clinical Assistant Professor, Quinnipiac University, Hamden CT, USA

Theodora Sirota, PhD, RN, CNL, PMHCNS-BC

Associate Professor, Department of Nursing, Caldwell College, Caldwell, NJ, USA

Diane Snow, PhD, RN, PMHNP-BC, CARN

Clinical Professor and Director, Psychiatric-Mental Health Nurse Practitioner Program, University of Texas at Arlington School of Nursing, Arlington, TX, USA

Roberta Waite, EdD, PMHCNS-BC, FAAN

Associate Professor and Assistant Dean of Academic Integration of Community Programs and Doctoral Nursing Department, Drexel University, Philadelphia,

PA, USA

Mary Weber, PhD, APRN, PMHNP-BC

Endowed Professor in Psychiatric Nursing, University of Colorado College of Nursing, Aurora, CO, USA

- Submit an Article
- Browse free sample issue
- **≜** Get content alerts
- Subscribe to this journal

More from this journal

News

Professional Opportunities

ORIGINAL ARTICLE



Determination of burnout, life satisfaction, and stress coping styles of parents with disabled children

Zehra Calıskan PhD. RN¹ Derva Evgin PhD. RN² Nuray Caner PhD. RN³



¹Child Health and Diseases Nursing Department, Nevsehir Hacı Bektaş Veli University, Nevsehir, Turkey

²Child Health and Diseases Nursing Department, Akdeniz University Kumluca Health Sciences Faculty, Antalya, Turkey

³Child Health and Diseases Nursing Department, Erciyes University, Kayseri,

⁴Child and Adolescent Psychiatry Department, Medical Faculty, İzmir Katip Çelebi University, İzmir, Türkiye

Correspondence

Nuray Caner, Child Health and Diseases Nursing Department, Health Sciences Faculty, Erciyes University, Kayseri 38260, Turkey. Email: goknuraytas@gmail.com

Funding information

Nevsehir Haci Bektas Veli University Scientific Research Projects, Grant/Award Number: NEÜBAP16SG38

Abstract

Purpose: This study determined problems, fatigue, life satisfaction, family stress, and coping strategies of families who had a disabled child and the relationships

Design and Methods: This descriptive and cross-sectional study conducted with 191 parents of children with disabilities.

Findings: As the life satisfaction of childcare providers increased, their emotional exhaustion decreased and sense of personal accomplishment increased (p < 0.05).

Practice Implications: It is necessary that nurses and health team members should provide regular and continuous education for burnout, life satisfaction, family stress and coping styles to parents of disabled children using interactive education techniques and group interactions.

KEYWORDS

burnout, coping with stress, disabled child, life satisfaction parent

1 | INTRODUCTION

Having a child with a disability is a crisis situation that can affect the quality of life of family members, the gender roles of parents, their economic expenses, their working status, their health, stress situations, their decision to maintain a marriage, and their decision to give birth to a new child.¹ The presence of a disabled person in the family generates ongoing problems in every period of life. Living with a disabled individual requires adaptation and causes some limitations in the lives of family members.^{2,3} The birth of the disabled child or the subsequent disability situation causes difficulties and changes as the family members adapt to the new conditions, family structure, and relationships with each other and their inner worlds.⁴ It is emphasized that this situation, which is difficult to get used to, affects parents' close environment economically, socially, and psychologically.^{2,3,5} In addition, disability in children affects

the emotional burden in the family, the social support for families is low,⁶ and the physical and mental health of mothers, especially, is negatively affected due to their higher responsibility for these children.⁷

Families need help and support in dealing with the stressful and long-term changes required to care for a child with disabilities. Therefore, it is important for nurses who care for children with disabilities and their families to know these families, to know their lives, and to plan family-centered nursing care. The nurse should make a holistic and family-centered care plan using many roles and independent functions such as educator, counselor, advocate, decision maker, and caregiver while carrying out her studies in this field.^{8,9} Research indicates that the gathering of families with similar problems has positive effects, professional support services for families reduces stress, family burdens,⁶ and burnouts, and increase their satisfaction. 10,11 Nurses, who have a responsibility to provide

This study was presented as an oral presentation in the 1st International 2nd National Public Health Nursing Congress, April 23-26, 2018, Ankara.

holistic care to families, should advise these families on coping with stress and take initiatives to increase their social support. ^{10,12,31} Despite descriptive studies in the literature regarding the problems experienced by families with disabled children, ^{1,13-18} we did not encounter any study emphasizing the relationship between burnout, life satisfaction, and coping styles of families with children with disabilities. Therefore, this study was needed.

2 | THE AIMS OF THE STUDY

This study determined problems, fatigue, life satisfaction, family stress, and coping strategies of families who had a disabled child and the relationships between them.

3 | METHODS

3.1 | Study design and population

This descriptive and cross-sectional study occurred in Turkey's centrally located province in Cappadocia, Family and Social Policies Provincial Directorate of Disability Services, located in the Department of Finance. Data was collected between April 2016 and June 2016. Parents of children with disabilities in the 0–18 age group were included.

The population of the study was composed of all parents who registered disabled children (*N*: 278) according to 2015 data. In the study, no sample calculations were made and all parents who voluntarily agreed to participate in the study were included. The research was completed with 191 parents.

4 | DATA COLLECTION TOOLS

4.1 | Information form

The researchers developed this based on the literature reviewed.^{6,19–21} The information form gathered the socio-demographic characteristics of the child with disabilities and their parents. Questions regarding the difficulties experienced by the parents in their individual and family life due to the disability in their children were included.

4.2 | Maslach Burnout Inventory

The scale, developed by Maslach and Jackson, consists of 22 items. The scale evaluates three subscales of burnout: "emotional exhaustion" (9 items), "personal accomplishment" (8 items), and "depersonalization" (5 items). From subscales of the Maslach Burnout Inventory, 0-36 points can be obtained for the emotional exhaustion (Cronbach a = 0.83), 0-32 points can be obtained for the personal accomplishment (Cronbach a = 0.72) and 0-20 points can be obtained for the depersonalization (Cronbach a = 0.65)²² In this current study Cronbach a values of the

subscales were 0.84, 0.76, and 0.55, respectively. The scale does not give total points but gives the total score of each subscales. High burnout reflects a high score in the "emotional exhaustion" and "depersonalization" subscales and a low score in the "personal accomplishment" subscale. Moderate burnout reflects moderate scores for all three subscales. Low level burnout reflects low scores on the "emotional exhaustion" and "depersonalization" subscales and high scores on the "personal accomplishment" subscale. In scoring, three separate burnout points are calculated for each person.²²

4.3 Life Satisfaction Scale

On the scale developed by Diener in 1985, 23 participants are presented with five situations they are asked to score using a 7-grade scale as follows: (1) "I certainly agree" to (7) "I certainly do not agree." A minimum of 5 and a maximum of 35 total points can be obtained. According to the total score of the scale, high scores indicate higher life satisfaction. 24 In the Turkish validity and reliability study of the life satisfaction scale, the Cronbach a value is 0.88. in this study Cronbach a value of the Life satisfaction scale is 0.80.

4.4 | Stress Coping Styles Scale (SCSS)

This scale is a 30-item, 5-subscale scale developed by Folkman and Lazarus called Ways of Coping Inventory in 1980 to determine the ways individuals use to deal with general or pronounced stressful situations.²⁵ Siva carried out the first standardization study of the scale. The scale is a 4-point Likert type and self-assessing scale rated from "not suitable" to "very convenient." In the evaluation of the scale, the total score is not obtained, and separate scores are calculated for the subscales. The scale has two dimensions: problem-focused or emotion-focused. Effective approaches to problems are described as "self-confident," "optimistic," and "seeking social support" whereas ineffective approaches are "helpless" and "submissive." The range of scores that can be obtained from the subscales are 7–28 points for the self-confident (Cronbach a = 0.80), 5–25 points for the optimistic (Cronbach a = 0.68), 4–16 points for the seeking social support (Cronbach a = 0.47), 8–32 points for the helpless (Cronbach a = 0.73) and 6-24 points for the submissive (Cronbach a = 0.70). The score increase in the subscales is interpreted as the individuals use that method of coping more.²⁶ In this study Cronbach a values of the subscales were 0.79, 0.68, 0.50, 0.66, and 0.48, respectively. When categorizing ways of coping as effective and ineffective, internal consistencies were found to be 0.76 for effective and 0.69 for ineffective ways of coping.

4.5 | Ethics

Ethics committee approval from the University Non-Interventional Ethics Committee (2016/11.01) and written permission from the Provincial Directorate of Family and Social Policies, where the study

Perspectives in PSYCHIATRIC CARE—WILEY

was carried out, was obtained before starting the study. In addition, the participants were informed about the purpose of the study, were asked to participate in the study, and their verbal and written consent was obtained.

DATA ANALYSIS 5

The data were evaluated using a statistics program. Number, percentage, arithmetic mean, and standard deviation distribution were used to evaluate descriptive statistics. In addition, the data were evaluated using the significance test of the difference between the two means (independent t, Mann-Whitney U) and Pearson Correlation tests. The homogeneity of the variances was checked using the Levene test. Statistical significance of the results was evaluated at p < 0.05 level.

RESULTS

Looking at the sociodemographics of the parents, 60.7% of the mothers and 53.4% of the fathers were primary school graduates, the majority of the mothers (96.3%) did not work, and 79.1% of the fathers worked. Most families had a nuclear structure (77%), 39.8% had three children, and more than half of the parents (66.5%) stated their income status as "less income than expenses" (Table 1).

It was determined that 60.7% of the children were male, 46.6% were in the 13-18 age group, and 67% went to the rehabilitation center. Most (68.1%) children were born vaginally, 57.1% had disabilities that emerged after birth, and 33.5% had more than one type of disability. The problems experienced from their disability include speech problems (51.9%) and bed sores (3.8%) (Table 2).

The majority of the families (86.9%) had difficulties in caring for the disabled child in general and most of them had difficulties in meeting hygene requirements (61.3%) as well as communicating, education, walking, dressing, and eating (60.3%, 59.7%, 57.1%, 52.4%, and 48.7%, respectively) (Table 3).

Parents whose children were in the 0-5 age group used the submissive approach more (p < 0.05). There was no statistically significant difference between the gender of the child and the parents' SCSS subgroup scores (for all: p > 0.05). While 57% of the parents stated they thought they were not understood by the people around them, 61.1% stated that they felt "sadness, disappointment, anxiety" for their disabled children. When questioned about the impact of their child's disability diagnosis on their family and social relationships, the relationships of parents with healthy children, relatives, neighbors, and their spouses (53.9%, 48.7%, 48.2%, and 41.9%, respectively) were affected and the majority (89.5%) experienced financial difficulties.

Parents who received support in the care of the disabled child used the social support behavior and parents who do not receive support in the care of the child used the submissive approach (Table 4) (p < 0.05). Parents who stated that they had difficulty

TABLE 1 Socio-demographics of parents

Family characteristics	Count	%
Responding person		
Mother	169	88.4
Father	11	5.8
Other ^a	11	5.8
Mother's education status		
Not literate	29	15.2
Primary school graduate	116	60.7
Middle school graduate	15	7.9
High school graduate	31	13.1
University graduate	2	1.0
No answer	4	2.1
Father's education status		
Not literate	9	4.7
Primary school graduate	102	53.4
Middle school graduate	33	17.3
High school graduate	34	17.8
University graduate	6	3.1
No answer	7	3.7
Mother working status		
Working	4	2.1
Not working	184	96.3
No answer	3	1.6
Father working status		
Working	151	79.1
Not working	29	15.2
No answer	10	5.2
Family structure		
Nuclear family	147	77.0
Extended family	33	17.3
Broken family	11	5.7
Perceived income status		
Revenue is less than expenses	127	66.5
Revenue is more than expenses	11	5.8
Revenue is equal to expenses	53	27.7
Number of children in the family		
One	14	7.3
Two	61	31.9
Three	76	39.8
Four or more	40	21.0

^aOther: grandmother and Grandfather.

meeting the hygiene requirements of the child used the desperate and submissive approach (p < 0.05). Parents who stated they had difficulties in getting the child around, eating, and dressing also used the submissive approach (p < 0.05) (Table 4).

Those with no support, including from their spouse, had less life satisfaction (p < 0.05). In addition, those who stated that they had difficulty in all kinds of care (walking, communication, dressing,

TABLE 2 Characteristics of children with disabilities

Characteristics of children with disabilities	Count	%
Gender Female Male	75 116	39.3 60.7
Age 0-3 4-6 7-12 13-18	10 22 70 89	5.1 11.6 36.7 46.6
Going to the rehabilitation center Going Not going	128 63	67.0 33.0
Form of delivery Normal vaginal delivery Cesarean delivery	130 61	68.1 31.9
Time disability was identified At birth After birth	82 109	42.9 57.1
Disability type Physically disabled Mildly mentally disabled Moderately mentally disabled Severely mentally disabled More than one type of disability Other ^a	28 8 39 48 64	14.7 4.2 20.4 25.1 33.5 2.1
Problems faced due to the disability (n: 264) ^b Hearing Talking Walking Bedsores	21 137 96 10	7.9 51.9 36.4 3.8

^aOther: deaf, visually impaired.

education, and feeding) had less life satisfaction (p < 0.05) and more emotional exhaustion (p < 0.05) (Table 5).

There was a moderately negative relationship between life satisfaction and emotional exhaustion of the caregivers of the disabled child and a weakly positive relationship between life satisfaction and personal accomplishment (p < 0.01) (Table 6).

7 | DISCUSSION

The age of the disabled child can influence the difficulties experienced by the mothers and their coping strategies. According to the study of Oymak and Arslan, ¹⁴ mothers of young children experience more problems than those of older children. One of the important findings in this study is that parents whose

TABLE 3 Fields where families have difficulty in caring for a disabled child

disabled child		
Difficulties experienced	Count	%
General Having difficulties Not having difficulties	166 25	86.9 13.1
Walking Having difficulties Not having difficulties No answer	109 80 2	57.1 41.9 1.0
Communication Having difficulties Not having difficulties No answer	115 74 2	60.3 38.7 1.0
Meeting the hygiene requirement Having difficulties Not having difficulties No answer	117 72 2	61.3 37.7 1.0
Dressing Having difficulties Not having difficulties No answer	100 89 2	52.4 46.6 1.0
Feeding Having difficulties Not having difficulties No answer	93 96 2	48.7 50.3 1.0
Education Having difficulties Not having difficulties No answer	114 75 2	59.7 39.3 1.0
Supporting person(s) Present Absent	103 88	53.9 46.1
Supporting person(s) ^a My family My friends My partner My children Reeve, municipality	73 5 28 8 1	38.2 2.6 14.7 4.1 0.5
Being understood by other people Yes No	82 109	43.0 57.0
Feelings as a mother of a disabled child ^a "I see him/her as a gift given to me, I am thankful" "Sadness, astringency, anxiety" "I got used to it" "It is very difficult, you cannot know unless you live with it"	15 118 15 17	7.8 61.1 7.8 8.8
Other ^b	28	14.5

^bMore than one answer was given. Percentages are calculated on "n" values.

TABLE 3 (Continued)

Difficulties experienced	Count	%
Unanswered	7	3.1

Note: The bold values indicate the percentage of the most common responses given by parents.

^bOther ("I am hopeful, I think it is a test, I feel tired, I cannot get into the community, I am overwhelmed, I am hopeless...").

children were in the 0-5 age group used more of the submissive approach to cope with stress (p < 0.05) (Table 4). If the disabled child is both in the young age group and has more than one disability, they have increased dependence on the parent caregiver. In this study, 86.9% of the participants stated that had difficulty caring for children with disabilities, primarily in communicating, educating, walking, dressing, and eating (60.3%, 59.7%, 57.1%, 52.4%, and 48.7%, respectively) (Table 3). Parents who had these difficulties used the submissive approach to cope with stress (p < 0.05) (Table 4). These results show it is difficult for mothers to care for their disabled child and they use ineffective methods for coping with these difficulties. Masulani-Mwale et al. 19 stated it is tiring for the mothers when the children are dependent on them, but mothers feel compelled to fill this role. The mother often takes full responsibility for her child, does not have sufficient support.²⁷ especially in the nuclear family, and has difficulties performing other roles.²⁸ At this point, parents need professional and social support.²⁹ The reason mothers need support is that the rate of mothers' participation in childcare is higher than that of fathers. 10 In the study, mothers who received support from their spouses in the care of the disabled child were low (14.7%). This is an important finding because it shows the mother plays the greatest role in the care of the disabled child, and sometimes is the only caregiver. Similar to the findings of this study, others have found the father's support in the care of the disabled child is low. 30 Because childcare and housework are considered the primary duty of women in a patriarchal society, this situation increases the burden of the mother in the care of the disabled child.

The disability often changes the daily routines of caregivers due to the workload it brings and can often lead to negative psychological and social effects. While 57% of the mothers stated that they thought that they were not understood by the people around them, 61.1% stated they felt felt "sadness, disappointment, anxiety" for their disabled children. In another study, parents felt sad after learning about the discomfort of their children, but most of these parents eventually accepted the situation due to their belief in fate. Some parents with sadness could not accept the seriousness of the situation and the negative perspective of society. In the study, after their children were diagnosed as "disabled," their family and social

relationships were negatively affected and most of them experienced financial problems. In the study conducted by Kahriman et al.,³² the relationships of individuals with disabled children with their spouses, friends, relatives, and neighbors were negatively affected. Having a child with a disability is a phenomenon that is difficult to accept for the family, restricting and complicating their individual and social lives.³¹

Mothers who have disabled children have difficulties in the care and treatment of their children and they experience burnout, hopelessness, depression, and stress. ^{13,15,17,18,30} The stress that families experience not only affects their ability to care for the child, but also affects their mental and physical health and the quality of their relationship with their children. ²¹ Therefore, the coping styles used by the parents to handle this stress are important for well-being and psychological health.

In this study, the parents who received support in the care of the disabled child used the social support behavior, and the parents who did not receive child care support used the submissive approach (p < 0.05) (Table 4). Similarly in another study, it was found that disabled childrens' mothers who are received social support, use effective ways of coping with stress.¹⁴ The supportive attitudes of relatives, close friends, and spouses, which are described as social support, are an important factor in coping.^{10,21}

When tackling a problem-oriented coping strategy, there is a tendency to actively solve the problem or passive coping, such as avoidance, depersonalization, and ignoring. When using a problem-oriented coping strategy, the problem is actively solved by seeking social support and using an optimistic and self-confident approach. However, using emotional-oriented coping relies on avoiding, depersonalization, and submissive approaches. Although positive emotional coping tactics are necessary for the individual to maintain his psychological well-being, long-term passive tactics do not solve the problem.²⁵

In this study, the life satisfaction of those who received support and those who received support from their husbands in coping with the problems they experienced in caring for disabled children were higher (p < 0.05). Community relations and family relations affect the family's quality of life positively.³³

Parents may find it difficult to cope with the problems (physical, emotional, and social) that the child experiences. After addressing the physical problems and the increase in workload, care givers may experience a sense of exhaustion if professional support is not received. In this current study, emotional exhaustion scores of mothers who stated that they had difficulties in all kinds of care of children with disabilities were higher (p < 0.05). The burnout level increases as the severity of the child's disability increases. Families need support systems to deal with the problems faced by families with disabled children, especially the high level of stress and the burnout that occurs after the more intense difficulties. Total Since intense emotional burden causes serious psychosocial problems in mothers, it is

^aMore than one answer was given.

TABLE 4 Subgroup point averages of the families receiving support in the care of the disabled child, the areas where they had difficulties, and the SCSS

and the SCSS					
	SCSS subgroups				
	Confident approach	Optimistic approach Median	Desperate approach	Submissive approach Median	Searching for social support Median
Problems experienced	(25th-75th percentile)	(25th-75th percentile)	(25th-75th percentile)	(25th-75th percentile)	(25th-75th percentile)
Age of child					
0-5 years	14.50	10.00	13.00 ± 4.35	11.06 ± 1.89	6.50
	(12.75-19.00)	(9.75–12.00)	(5.00-22.00)	(8.00-14.00)	(6.00-8.00)
6-12 years	15.00	11.00	11.67 ± 3.99	9.35 ± 2.92	7.00
12 10	(14.00–19.00)	(10.00-13.00)	(3.00-21.00)	(3.00-18.00)	(6.00-9.00)
13-18 years	16.00 (14.00-19.00)	11.00 (10.00-13.00)	11.61 ± 4.67 (2.00-24.00)	9.72 ± 2.58 (4.00–15.00)	7.00 (6.00–9.00)
	$\chi^2 = 2.268$	$\chi^2 = 0.639$	(2.00-24.00) F = 0.873	(4.00–15.00) F = 3.093	$\chi^2 = 2.388$
	$\chi = 2.200$ df = 2	$\chi = 0.037$ $df = 2$	p = 0.420	p = 0.048	$\chi = 2.566$ df = 2
	p = 0.322	p = 0.726	ρ 0.420	p 0.040	p = 0.303
Compart in shilds are	r	F			F
Support in childcare Present	16.00	11.00	12.00	9.31 ± 2.67	8.00
Tresent	(14.00-20.00)	(10.00-13.00)	(9.00-14.00)	(3.00-15.00)	(6.00-9.00)
Absent	15.00	10.00	12.00	10.23 ± 2.69	7.00
	(14.00-18.00)	10.00-12.00)	(9.00-14.00)	(3.00-18.00)	(5.00-8.00)
	U = 3723.00	<i>U</i> = 3840.00	U = 4332.00	t = -2.342	U = 3286.00
	p = 0.072	p = 0.135	p = 0.884	p = 0.020	p = 0.003
Difficulty traveling wit	h the child				
Experiencing	16.00	11.00	12.59 ± 4.37	10.00	7.66 ± 2.08
	(14.00-19.00)	(10.00-14.00)	(2.00-22.00	(8.00-12.00)	(3.00-12.00)
Not experiencing	16.00	11.00	11.19 ± 4.11	9.00	7.70 ± 2.24
	(14.00-20.00)	(10.00-12.00)	(5.00-21.00)	(7.00-10.00)	(1.00-12.00)
	<i>U</i> = 3660.00	<i>U</i> = 4129.00	t = 1.863	U = 3195.00	t = −0.827
	p = 0.095	p = 0.703	p = 0.064	p = 0.003	p = 0.409
Difficulty in meeting the	he hygiene requirement of t	he child			
Experiencing	16.00	11.00	12.51 ± 4.26	9.69 ± 2.46	7.00
	(14.00-18.00)	(10.00-13.00)	(2.00-22.00)	(4.00–15.00)	(6.00-9.00)
Not experiencing	17.50	11.00	11.20 ± 4.28	8.83 ± 2.90	8.00
	(14.00–20.00) U = 3580.00	(10.00–14.00) U = 3764.00	(3.00–21.00) t = 2.277	(3.00–15.00) t = 2.524	(6.00-9.00) U = 3596.50
	p = 0.148	p = 0.347	p = 0.024	p = 0.012	p = 0.159
Difficulty in dressing to	•	F 313 11	F	F	F 3.23.
Experiencing	15.00	11.00	12.87 ± 4.36	9.98 ± 2.50	7.49 ± 2.40
Experiencing	(14.00-19.00)	(10.00-13.00)	(2.00-22.00)	(3.00-15.00)	(1.00-12.00)
Not Experiencing	17.00	11.50	11.02 ± 4.04	8.65 ± 2.69	7.88 ± 1.82
, ,	(14.00-20.00)	(10.00-13.75)	(2.00-21.00)	(4.00-15.00)	(3.00-12.00)
	<i>U</i> = 3711.50	<i>U</i> = 3899.00	t = 1.206	t = 2.847	t = -1.042
	p = 0.081	p = 0.214	p = 0.229	p = 0.005	p = 0.299
Difficulty in feeding th	e child				
Experiencing	15.00	10.50	12.87 ± 4.22	10.09 ± 2.38	7.41 ± 2.28
	(14.00-18.00)	(9.00-13.00)	(2.00-22.00)	(4.00-15.00)	(1.00-12.00)
Not Experiencing	17.00	12.00	11.25 ± 4.25	8.73 ± 2.75	7.89 ± 2.02
	(14.00-20.00)	(10.00-14.00)	(2.00-21.00)	(3.00-15.00)	(3.00-12.00)
	U = 3606.50	U = 3538.00	t = 1.284	t = 2.675	t = −1.793
	p = 0.038	p = 0.022	p = 0.201	p = 0.008	p = 0.076

Abbreviations: F, one way ANOVA test; SCSS, Stress Coping Styles Scale; t, independent T test; U, Mann-Whitney U test.

TABLE 5 Subgroup averages of the parents, the areas where they have difficulties in the care of the child, and the Life Satisfaction and Maslach Burnout Inventory

Variables	Life satisfaction Median (25-75	Personal accomplishment Median	Emotional exhaustion Median (25-75	Desensitization Median
Comment was a few	percentile)	(25-75 percentile)	percentile)	(25-75 percentile)
Support person for	•	07.0 (05.0, 00.0)	40.0 ((0.45.0)	0.0 (0.0 5.0)
Present Absent	19.0 (15.0–25.0) 15.0 (11.75–20.25)	27.0 (25.0–30.0) 26.0 (21.0–29.0)	10.0 (6.0–15.0) 12.5 (6.0–18.0)	3.0 (0.0–5.0) 3.5 (2.0–5.0)
	U = 3249.000 p = 0.002	U = 37.71.500 p = 0.078	U = 38.24.000 p = 0.106	U = 4186.000 p = 0.511
Spouse support in	the care of the child			
Present	18.0 (13.0-24.25)	27.0 (23.75-30.0)	11.0 (6.0-16.0)	3.0 (1.0-5.0)
Absent	15.0 (10.25-20.0)	26.0 (21.5-28.0)	10.5 (7.0-19.0)	3.0 (0.0-7.0)
	U = 2577.500	U = 3153.000	<i>U</i> = 3060.000	U = 3390.500
	p = 0.012	p = 0.437	p = 0.293	p = 0.957
Difficulty in the ge	neral care of the chil	d		
Experiencing	17.0 (12.0-22.0)	26.0 (22.75-29.0)	12.0 (7.0-17.0)	3.0 (0.0-5.0)
Not experiencing	25.0 (17.5-29.0)	28.0 (25.5-31.0)	6.0 (2.0-10.5)	3.0 (0.5-5.0)
	U = 1225.000	<i>U</i> = 1646.500	U = 1106.500	<i>U</i> = 1994.000
	p = 0.001	p = 0.121	p < 0.001	p = 0.750
Difficulty traveling	with the child			
Experiencing	17.0 (11.0-22.0)	26.0 (22.0-29.0)	13.0 (8.08-18.0)	3.0 (0.25-5.75)
Not experiencing	19.5 (14.0-25.75)	27.0 (25.0-30.0)	8.5 (4.5-14.75)	3.0 (0.0-5.0)
	U = 3255.000	U = 3758.500	U = 2912.000	U = 4246.500
	p = 0.004	p = 0.127	p < 0.001	p = 0.840
Difficulty in comm	unicating with the ch	ild		
Experiencing	17.0 (12.0-22.0)	26.0 (22.0-28.0)	13.0 (7.0-17.0)	4.0 (1.0-6.0)
Not experiencing	19.0 (13.75-26.0)	27.5 (24.0-30.0)	9.5 (5.0-14.0)	2.5 (0.0-5.0)
	U = 3414.500	U = 3465.500	U = 3356.500	U = 3739.500
	p = 0.022	p = 0.031	p = 0.014	p = 0.154
Difficulty in meeting	ng the hygiene require	ement of the child		
Experiencing	17.0 (12.5-22.0)	26.0 (23.0-29.0)	12.0 (7.0-17.0)	3.0 (0.0-5.0)
Not experiencing	19.0 (12.25-26.0)	27.0 (22.5-30.0)	9.5 (5.25-15.0)	3.0 (1.0-5.0)
	<i>U</i> = 3489.000	<i>U</i> = 3717.500	<i>U</i> = 3416.000	<i>U</i> = 4160.500
	p = 0.048	p = 0.174	p = 0.029	p = 0.886
Difficulty in dressin	ng the child			
Experiencing	16.0 (11.0-21.75)	26.0 (22.25-29.0)	14.0 (9.0-17.75)	4.0 (1.0-6.0)
Not experiencing	19.0 (15.0-24.5)	27.0 (24.0-30.0)	8.0 (4.5-14.0)	2.0 (0.0-4.5)
	U = 3387.000	U = 4003.500	U = 2972.500	U = 3694.000
	p = 0.005	p = 0.233	p < 0.001	p = 0.041
Difficulty in the ed	lucation of the child			
Experiencing	17.0 (12.0-22.0)	26.0 (22.0-29.0)	13.0 (8.0-18.0)	4.0 (1.0-6.0)
Not experiencing		27.0 (24.0-30.0)	8.0 (4.0-15.0)	2.0 (0.0-4.0)
	<i>U</i> = 3472.000	<i>U</i> = 3906.500	<i>U</i> = 3030.000	U = 3629.500
Diff. It is a	p = 0.029	p = 0.315	p = 0.001	p = 0.075
Difficulty in feeding		0.40.45.45.55.55		
Experiencing	17.0 (12.0-22.0)	26.0 (21.0-28.0)	14.0 (.0-18.5)	4.0 (1.0-5.5)
Not experiencing	19.0 (13.0-25.0)	27.5 (24.25–30.0)	9.0 (4.0-14.0)	2.0 (0.0-5.0)
	<i>U</i> = 3519.000	<i>U</i> = 3555.000	U = 2995.000	<i>U</i> = 4074.500
	p = 0.012	p = 0.015	p < .001	p = .294

TABLE 6 Correlation between parents' life satisfaction and Maslach Burnout Inventory Subgroup Mean Scores

	Life satisfaction	Personal accomplishment	Emotional exhaustion	Desensitization
	r	r	r	r
Life satisfaction	1.000			
Personal accomplishment	0.328*	1.000		
Emotional exhaustion	-0.465*	-0.420*	1.000	
Desensitization	-0.277*	-0.315	0.493*	
				1.000

Abbreviations: r, Pearson correlation coefficient.

*p < 0.01.

Pearson r

0 < r < 0.20: very weak correlation.

 $0.20 \le r < 0.40$: Weak correlation.

 $0.40 \le r \le 0.60$: Moderate correlation.

 $0.60 \le r < 0.80$: Good correlation.

 $0.80 \le r \le 1$: Strong correlation.

important to address the difficulties experienced by mothers professionally. Education and counseling for the family can reduce the high level of stress and burnout experienced, especially for the mother.

The most important result of the study is that as the life satisfaction of those who care for the disabled child increases, their personal accomplishment increases (p < 0.05) and their emotional exhaustion decreases (p < 0.01). Similar to this study, a negative correlation was found between the burnout level of the mothers who have disabled children and their perceptions of quality of life and life satisfaction. Social and educational support for families with children with disabilities can contribute to increasing life satisfaction and personal achievement levels and reducing burnout levels. Nurses can try to reduce the burden and stress of the family and help them cope with it through efforts intended to meet the needs of families with children with disabilities. In addition to these activities, nurses should take a more active role in the education and support provided to the families of children with mental disabilities. 10

8 | CONCLUSION

Having a disabled child can affect parents' gender roles, economic expenses, health, quality of life and satisfaction, stress levels and coping with stress abilities. Parents' getting adequate social support and getting professional help both increase their life satisfaction and enable them to use effective methods of coping with stress. The most important result of the study was that as the life satisfaction of those who care for the disabled child increases, their personal accomplishment increases and their emotional exhaustion decreases. According to the results obtained, it is recommended that:

 Regular monitoring of families with children with disabilities between the ages of 0-5, which is a high risk group for stress mismanagement, and establishing social support groups,

- In the Provincial Directorates of Family and Social Policies, regular and continuous training is provided to the mothers of children with disabilities, using interactive teaching techniques and providing effective group interactions, by a professional team, including nurses,
- Planning the interventions according to the levels of families' burnout, life satisfaction, and ways of dealing with stress at regular intervals after the children are diagnosed with a disability,
- Community trainings to change the negative views of the society towards children with disabilities and to accept these children.

9 | IMPLICATIONS FOR PSYCHIATRIC NURSING PRACTICE

Having a disabled child can affect parents' gender roles, economic expenses, health, quality of life and satisfaction, stress and coping with stress, and burnout levels, depending on the difficulties experienced in the care of the child. Parents' having adequate social support and professional help both increase their life satisfaction and enable them to use effective methods of coping with stress. The most important result of the study: as the life satisfaction increases, their personal success increases and their emotional exhaustion decreases of caregivers of disabled children. It is important for nurses who care for disabled children and their families in clinics, rehabilitation centers and the society in terms of getting to know these families, knowing their lives, and planning nursing care.

10 | LIMITATIONS OF THE STUDY

The collection of data mostly from mothers is a limitation of the research. More fathers' answers to questions may lead to differences in results. For this reason, it is recommended that both mothers' and

fathers' satisfaction with life, coping with stress, and burnout levels should be examined and compared in future research. In addition, how the siblings of mentally handicapped children are affected should be examined.

ACKNOWLEDGEMENTS

This study was supported by Nevşehir Hacı Bektaş Veli University Scientific Research Projects (Project No: NEÜBAP16SĞ38). The Project name is "The Effects of Education for Mothers with Disabilities on Burnout, Life Satisfaction, Family Stress and Coping Styles of Mothers." The Project manager was Zehra Çalışkan. Derya EVGİN, Nuray CANER, Bahriye KAPLAN, and Gonca ÖZYURT worked in the project as researchers. The Project started on 01 APRÌL 2016, and finished on 29 December 2017.

CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

AUTHOR CONTRIBUTIONS

Concept: Zehra Çalışkan. Design: Zehra Çalışkan and Derya Evgin. Supervision: Zehra Çalışkan and Derya Evgin. Financing: Zehra Çalışkan, Derya Evgin, Nuray Caner, and Bahriye Kaplan. Material: Zehra Çalışkan. Data collection or processing: Zehra Çalışkan, Bahriye Kaplan, Nuray Caner, and Gonca Özyurt. Analysis and interpretation: Zehra Çalışkan, Derya Evgin, Nuray Caner, Bahriye Kaplan, and Gonca Özyurt. Literature search: Zehra Çalışkan and Derya Evgin. Posted by: Derya Evgin and Zehra Çalışkan. Critical revision: Zehra Çalışkan.

ORCID

Zehra Çalışkan https://orcid.org/0000-0002-4726-5052

Derya Evgin https://orcid.org/0000-0002-3452-2937

Nuray Caner https://orcid.org/0000-0001-7817-7250

Bahriye Kaplan https://orcid.org/0000-0003-1600-3188

Gonca Özyurt https://orcid.org/0000-0002-0508-0594

REFERENCES

- Giulio PD, Philipov D, Jaschinski I. Families with disabled children in different european countries. Fam Soc. 2014;23:1-47.
- Lima-Rodríguez JS, Baena-Ariza MT, Domínguez-Sánchez I, Lima-Serrano M. Intellectual disability in children and teenagers: influence on family and family health. Systematic review. Enfermería Clínica (English Edition). 2018;28(2):89-102. https://doi.org/10.1016/j.enfcli.2017.10.005
- Van Cleave J. Children with special health care needs: with population-based data, better individual care plans. *Pediatrics*. 2015; 135(4):e1040-e1041. https://doi.org/10.1542/peds.2015-0319
- Rayan A, Ahmad M. Mindfulness and parenting distress among parents of children with disabilities: a literature review. *Perspect Psychiatr Care*. 2018;54(2):324-330. https://doi.org/10.1111/ppc. 12217
- Whiting M. What it means to be the parent of a child with a disability or complex health need. Nurs Child Young People. 2014;26(5): 26-29. https://doi.org/10.7748/ncyp.26.5.26.e390
- Caliskan Z, Bayat M. The effect of education and group interaction on the family burden and support in the mothers of intellectually

- disabled children. *Anatolian J Psychiatry*. 2016;17(3):214-223. https://doi.org/10.5455/apd.184412
- Sadati AK, Salehzade H, Hemmati S, Darvish M, Heydari ST, Tabrizi R. The causal factors associated with the loving care of the mothers of children with multiple disabilities. *Int J Community Based Nurs Midwifery*. 2015;3(4):309.
- Almasri NA, An M, Palisano RJ. Parents' perception of receiving family-centered care for their children with physical disabilities: a meta-analysis. *Phys Occup Ther Pediatr*. 2018;38(4):427-443. https://doi.org/10.1080/01942638.2017.1337664
- Chambers HG, Chambers JA. Effects of caregiving on the families of children and adults with disabilities. *Phys Med Rehabil Clin*. 2015; 26(1):1-19. https://doi.org/10.1016/j.pmr.2014.09.004
- inan Budak M, Küçük L, Civelek HY. Life experiences of mothers of children with an intellectual disability: a qualitative study. J Mental Health Res Intellectual Disabil. 2018;11(4):301-321. https://doi.org/ 10.1080/19315864.2018.1518502
- Sohmaran C, Shorey S. Psychological interventions in reducing stress, depression and anxiety among parents of children and adolescents with developmental disabilities: a systematic review and meta-analysis. J Adv Nurs. 2019;75(12):3316-3330. https://doi.org/ 10.1111/jan.14166
- Coren E, Ramsbotham K, Gschwandtner M. Parent training interventions for parents with intellectual disability. *Cochrane Database Syst Rev.* 2018;137(7):CD007987. https://doi.org/10.1002/14651858.CD007987
- Aktan O, Orakcı Ş, Durnalı M. Investigation of the relationship between burnout, life satisfaction and quality of life in parents of children with disabilities. Eur J Special Needs Education. 2020;35:1-17. https://doi.org/10.1080/08856257.2020.1748429
- Oymak HA, Arslan FT. Ways of coping with stress and perceived social support among mothers of children with intellectual disability.
 J Contemp Med. 2019;10:1-7. https://doi.org/10.16899/jcm.739169
- Sekułowicz M, Boroń-Krupińska K, Kwiatkowski P, Latawiec BM. Sense of coherence, resilience and coping as correlates of burnout of parents of children with disabilities: the new burnout screening instrument. Niepełnosprawność-Dyskursy Pedagogiki Specjalnej. 2019; 36:254-268.
- Yilmaz G. Spiritual orientation, meaning in life, life satisfaction, and well-being in mothers with disabled children. J Relig Health. 2019; 58(6):2251-2262. https://doi.org/10.1007/s10943-019-00925-4
- Yilmaz G. Mothers with disabled children: needs, stress levels and family functionality in rehabilitation. *Scand J Caring Sci.* 2020;34(2): 524-532. https://doi.org/10.1111/scs.12783
- Yilmaz G, Küçük Alemdar D. Evaluation of care burden among mothers of children with a disability: correlation between physical activity, quality of life, and sleep quality: a cross-sectional study. Perspect Psychiatr Care. 2020;57:129-137. https://doi.org/10.1111/ ppc.12534
- Masulani-Mwale C, Mathanga D, Silungwe D, Kauye F, Gladstone M. Parenting children with intellectual disabilities in Malawi: the impact that reaches beyond coping? *Child Care Health Dev.* 2016;42(6): 871-880. https://doi.org/10.1111/cch.12368
- Mukherjee M, Shignapure V. Challenges faced by parent due to the presence of mentally handicapped person in the family. *Int J Indian Psychol.* 2016;3(3):61-77.
- Peer JW, Hillman SB. Stress and resilience for parents of children with intellectual and developmental disabilities: a review of key factors and recommendations for practitioners. J Policy Pract Intellectual Disabil. 2014;11(2):92-98. https://doi.org/10.1111/jppi. 12072
- 22. Çapri B. Turkish adaptation of the burnout measure: a reliability and validity study. *Mersin Uni J Faculty Edu*. 2006;2(1):62-77.
- Diener E, Emmons RA, Larsen RJ, Griffin S. The satisfaction with life scale. J Pers Assess. 1985;49(1):71-75.

- Dağlı A, Baysal N. Adaptation of the satisfaction with life scale into Turkish: the study of validity and reliability. *Electronic J Soc Sci.* 2016;15(59):1250-1262.
- 25. Şahin NH, Durak A. Stress-coping styles scale: adapted for university students. *Turkish J Psychol.* 1995;10(34):56-73.
- Evgin D, Erdem E. The ways of coping stresss and levels of depression of the parents of children with chronic renal failure. Bozok Med J. 2018;8(2):33-41.
- Sato N, Araki A, Ito R, Ishigaki K. Exploring the beliefs of Japanese mothers caring for a child with disabilities. *J Fam Nurs*. 2015;21(2): 232-260. https://doi.org/10.1177/1074840715586551
- Čagran B, Schmidt M, Brown I. Assessment of the quality of life in families with children who have intellectual and developmental disabilities in Slovenia. J Intellectual Disabil Res. 2011;55(12): 1164-1175. https://doi.org/10.1111/j.1365-2788.2011.01400.x
- Hamedanchi A, Khankeh HR, Fadayevatan R, Teymouri R, Sahaf R. Bitter experiences of elderly parents of children with intellectual disabilities: a phenomenological study. *Iran J Nurs Midwifery Res.* 2016;21(3):278. https://doi.org/10.4103/1735-9066.180385
- Bawalsah JA. Stress and coping strategies in parents of children with physical, mental, and hearing disabilities in Jordan. *Int J Edu*. 2016;8(1):1-22. https://doi.org/10.5296/ije.v8i1.8811

- Crnic KA, Neece CL, McIntyre LL, Blacher J, Baker BL. Intellectual disability and developmental risk: promoting intervention to improve child and family well-being. *Child Dev.* 2017;88(2):436-445. https://doi.org/10.1111/cdev.12740
- Kahriman İ, Polat S, Gürol A. Determination of perceived social support and burnout levels of mothers of children with intellectual disability. J Pediatr Res. 2019;6(4):266-280. https://doi.org/10.4274/ jpr.galenos.2019.86648
- Schmidt J, Schmidt M, Brown I. Quality of life among families of children with intellectual disabilities: a Slovene study. J Policy Pract Intellectual Disabil. 2016;14(1):87-102. https://doi.org/10.1111/jppi. 12188

How to cite this article: Çalişkan Z, Evgin D, Caner N, Kaplan B, Özyurt G. Determination of burnout, life satisfaction, and stress coping styles of parents with disabled children. *Perspect Psychiatr Care*. 2021;57: 1505–1514. https://doi.org/10.1111/ppc.12746